

NOTES ON NEW AND NOTEWORTHY PLANTS. CXVII

Harold N. Moldenke

PHYLA NODIFLORA var. ANTILLANA Moldenke, var. nov.

Haec varietas a forma typica speciei recedit laminis foliorum margine argutissime patenteque dentatis, dentibus perconspicuis anguste triangularibus peracutis late divaricatis vel subreflexis.

This variety differs from the typical form and all other named forms and varieties of this species in having the margins of its leaf-blades very conspicuously spreading-dentate, the teeth large, narrowly triangular, sharply acute apically, and widely divaricate or even subreflexed.

The type of the variety was collected by Nathaniel Lord Britton, Elizabeth Gertrude Britton, and James F. Kemp (no. 83) on a hillside at Judith's Fancy, St. Croix, Virgin Islands, between March 17 and 25, 1923, and is deposited in the Britton Herbarium at the New York Botanical Garden.

VERBENA SIMPLEX f. EGGERTI (Moldenke) Moldenke, stat. nov.

Verbena simplex var. eggerti Moldenke, Am. Midl. Nat. 24: 753. 1940.

ADDITIONAL MATERIALS TOWARD A MONOGRAPH OF THE GENUS CALLICARPA.
XXVIII

Harold N. Moldenke

CALLICARPA KOCHIANA f. VILLOSULA Moldenke, Phytologia 39: 424. 1978.

Bibliography: Moldenke, Phytologia 39: 424. 1978.

Citations: CHINA: Fukien: Lae-Kwan 31 (Mi--type).

CALLICARPA LANCIFOLIA Millsp.

Additional bibliography: Gillis, Phytologia 35: 94. 1976; Moldenke, Phytologia 34: 155, 253, 254, & 272. 1976.

Byrne refers to this plant as a "boarhog bush 2 m. tall, not very common, only 2 individuals seen; upper leaf-surface dark-green when fresh; used in local medicine as a tonic." The Corrells describe it as a straggly shrub with white flower-buds and encountered it at the edge of coppices, flowering in August.

Additional citations: BAHAMA ISLANDS: Cat: Byrne 279 (N). Euthera: Correll & Correll 48945 (N).